Preprint typeset in JHEP style - HYPER VERSION

arXiv:1304.2978v1 [hep-th] 10 Apr 2013

The Construction of Dual-trace Factor in Yang-Mills Theory

Yi-Jian Du [a], **Bo Feng** [b,c], **Chih-Hao Fu** [d]

[a] *Department of Physics and Center for Field Theory and Particle Physics, Fudan University, Shanghai 200433, P.R China*
[b] *Zhejiang Institute of Modern Physics, Zhejiang University, 38 Zheda Road Hangzhou, 310027 P.R China*
[c] *Center of Mathematical Science, Zhejiang University, 38 Zheda Road Hangzhou, 310027 P.R China*
[d] *Department of Electrophysics, National Chiao Tung University*
1001 University Road, Hsinchu, Taiwan, R.O.C.

`yjdu@fudan.edu.cn; b.feng@cms.zju.edu.cn; zhihaofu@nctu.edu.tw`

Abstract: Recently, it has been proposed that tree-level Yang-Mills amplitudes can be written in dual-trace form, where the dual-trace factors satisfy cyclic symmetry and KK-relation. In this paper, we present a systematic construction of dual-trace factors based on the proposed relations between them and kinematic numerators of BCJ form in DDM chain. We show that solutions from our algorithm satisfy the relabeling property. We have also shown that using the relabeling property, solutions can be obtained independently.

Keywords: Dual-color factor.

Contents

1. Introduction **1**

2. Useful properties of BCJ numerators **5**

3. Construction of τ by BCJ numerators **8**

4. Examples **10**
- 4.1 Three-point case 10
- 4.2 Four-point case 11
- 4.3 five-point case 12
- 4.4 Six-point case 14
- 4.5 Seven-point case 14

5. Relabeling property **15**
- 5.1 The proof of relabeling invariance 16
- 5.2 An application 18
- 5.3 The implication of permutation $P(1n)$ 21

6. Conclusion **22**

A. The relabeling of permutation P_{1i} **22**

1. Introduction

In recent years, a significant progress in the study of scattering amplitudes is the discovery of kinematic-color duality(Bern-Carrasco-Johansson duality[1]). For tree-level amplitudes, BCJ duality states that Yang-Mills tree amplitudes can always be written as the following formula[1]

$$\mathcal{A}_{tot} = \sum_i \frac{c_i n_i}{D_i}, \tag{1.1}$$

where the sum runs over all distinguished tree diagrams with only cubic vertexes. In this BCJ form, the kinematic factors n_i (we will call "BCJ numerator") satisfy the same algebraic identities as those of the color factors c_i, i.e.,

$$\text{antisymmetry :} \quad c_i \to -c_i \Rightarrow n_i \to -n_i$$

$$\text{Jacobi}-\text{like identity}: \quad c_i+c_j+c_k=0 \Rightarrow n_i+n_j+n_k=0. \tag{1.2}$$

The BCJ duality between kinematic factors and color factors provides a strong constraint on color-ordered tree-level amplitudes in Yang-Mills theory. More explicitly, the antisymmetry of kinematic factors implies Kleiss-Kuijf [2, 3] relation, while the Jacobi-like identity implies BCJ relation[1]. The newly discovered BCJ relation has been understood in both string theory [4, 5, 6] and field theory [7, 8, 9, 10]. BCJ relation also serves as the key to the understanding of KLT relation [11], which expresses gravity tree amplitudes in terms of products of two color-ordered Yang-Mills tree amplitudes (See [12, 13, 14, 15]). Although not yet proven, many evidences show that the BCJ duality (1.2) is also respected at loop levels [16, 17, 18, 19, 20, 21, 22, 23, 24, 25].

Since BCJ duality has many important applications in the study of amplitudes in Yang-Mills theory and gravity, the construction of BCJ numerators becomes also an important problem and there are many works for it. In [26], BCJ numerators are constructed by pure-spinor string method. In [27, 28], a light-cone gauge approach for kinematic algebra was suggested, which provides a natural algebraic explanation of BCJ duality. Using this approach numerators of Yang-Mills amplitude up to MHV and all amplitudes in self-dual Yang-Mills theory can indeed be expressed as structure constants of generators of a diffeomorphism algebra [27, 28]. Based on the above idea, in a previous paper we proposed a more general kinematic algebra [29], from which one can construct BCJ numerators at tree-level in arbitrary D-dimensions and for arbitrary helicity configuration.

Because the BCJ duality (1.2), the algebra played by kinematic numerator and algebra played by color have same structure, thus there are many possible ways to decompose tree-level Yang-Mills amplitudes. The two traditional color-decompositions in Yang-Mills theory are given by [3]

$$\text{Trace form}: \quad \mathcal{A}_{tot}=\sum_{\sigma\in S_{n-1}}\mathrm{Tr}(T^{\sigma_1}...T^{\sigma_n})A(\sigma) \tag{1.3}$$

$$\text{DDM form}: \quad \mathcal{A}_{tot}=\sum_{\sigma\in S_{n-2}}c_{1|\sigma(2,..,n-1)|n}A(1,\sigma,n) \tag{1.4}$$

where A are color ordered amplitudes, T^a is the matrix of fundamental representation of $U(N)$ group and $c_{1|\sigma(2,..,n-1)|n}$ are constructed using the structure constants f^{abc}. For example, we have

$$c_{1|\sigma(2,..,n-1)|n}=f^{\sigma_1\sigma_2x_1}f^{x_1\sigma_3x_2}...f^{x_{n-3}\sigma_{n-1}n} \tag{1.5}$$

The transformation from double-copy form (1.1) to DDM form was shown in [30] using the KLT relation of colored scalar theory, while the transformation from DDM form to Trace form was given in [3] where the following two properties of Lie algebra of $U(N)$ gauge group were essential

$$\text{Property One}: \quad (f^a)_{ij}=f^{aij}=\mathrm{Tr}(T^a[T^i,T^j]), \tag{1.6}$$

$$\text{Property Two}: \quad \sum_a\mathrm{Tr}(XT^a)\mathrm{Tr}(T^aY)=\mathrm{Tr}(XY) \tag{1.7}$$

Based on the BCJ duality (1.1), it is natural to exchange the role between c_i and n_i and consider the following two dual formulations

$$\text{Dual Trace form :} \qquad \mathcal{A}_{tot} = \sum_{\sigma \in S_{n-1}} \tau_{\sigma_1 \ldots \sigma_n} \widetilde{A}(\sigma) \tag{1.8}$$

$$\text{Dual DDM form :} \qquad \mathcal{A}_{tot} = \sum_{\sigma \in S_{n-2}} n_{1|\sigma(2,..,n-1)|n} \widetilde{A}(1,\sigma,n) \tag{1.9}$$

where $\widetilde{A}$ is color ordered tree amplitude of scalar theory with f^{abc} as the only cubic coupling constants (see the references [**?**, 30]) and τ (we will call it "Dual-trace factor" or simply τ-function) is required to be cyclic invariant. Indeed, the Dual-DDM was given in [31] while the Dual-Trace form was conjectured in [32] with explicit constructions for first few lower-point amplitudes and a general construction was suggested in [28].

Although the existence of above two dual formulations were established, a systematic Feynman rule-like prescription to the coefficients τ and n is not known at this moment. The dual DDM-form has been studied in [29], where the construction of numerator factor $n_{1|\sigma(2,..,n-1)|n}$ has been given (see (2.5)). Although results in [29] for BCJ numerators n are just a small step towards the systematic construction, it does give us some useful applications.

In this paper, we use BCJ numerators to systematically construct the dual-trace factor τ and realize the proposed Dual-trace formulation (1.8). Unlike the trace form where (1.6) and (1.7) are available, there is no any relation among these τ-functions, thus in principle there are $n!$ τ-functions we need to determine. Any solution to these $n!$ τ-functions will be a rightful choice as long as it gives the right total amplitude (1.8). However, from the dual-DDM form, we see that there are only $(n-2)!$ BCJ numerator factors n needed to completely fix the total amplitude, thus it is very natural to impose some relations among τ's to reduce the independent τ-functions. The imposed relations are [32]:

- (A) **Cyclic symmetry:**

  $$\tau_{12\ldots n} = \tau_{n1\ldots(n-1)}. \tag{1.10}$$

  Using the cyclic symmetry, we can fix the first index to be any particular number, for example, 1, thus the independent τ-functions will be $(n-1)!$.

- (B) **KK-relation:**

  $$\tau_{1,\alpha,n,\beta^T} = (-1)^{n_\beta} \sum_{\sigma \in OP(\{\alpha\} \bigcup \{\beta\})} \tau_{1,\sigma,n}, \tag{1.11}$$

  where n_β is the number of elements in the set β and β^T denotes the inverse of the ordering of set β. The sum in (1.11) is over all permutations of the set $\{\alpha\} \bigcup \{\beta\}$ where relative ordering inside each subsets α and β is kept. Using this relation, we can fix two particular numbers, for example, 1 and n, at the first and last positions, thus the number of independent τ-functions is $(n-2)!$.

Having imposed above two relations, we need to find these $(n-2)!$ independent τ-functions to satisfy relation (1.8). This problem is solved by imposing following relations between $(n-2)!$ BCJ numerator factors $n_{1\sigma n}$ (since the $1, n$ are fixed, to simplify notation, we will write $n_{1|\sigma|n}$ inside (1.9) to be $n_{1\sigma n}$) and $(n-2)!$ $\tau_{1\sigma n}$-functions (where $\sigma \in S_{n-2}(23...(n-1))$)

$$n_{12...(n-1)n} = \tau_{1[2,[...,[n-1,n]...]]}, \tag{1.12}$$

where [,] means the antisymmetric combinations. For example, $n_{123} = \tau_{1[2,3]} = \tau_{123} - \tau_{132}$.

After solving $\tau_{1\sigma n}$ as linear combinations of $n_{1\sigma n}$ using (1.12), we can use (1.11) and (1.10) to obtain all other τ-functions. The claim is that if we put these τ's back to the right handed side of (1.8), then we do get the left handed side. The key to guarantee the claim is relations (1.12). In fact, the proof of equivalence of the trace form (1.3) and the DDM form (1.4) based only following two facts: (1) partial amplitudes $A(1, \sigma, n)$ are cyclic and satisfy KK-relations; (2) two different color factors satisfy

$$c_{12...(n-1)n} = \mathrm{Tr}(T^{a_1}[T^{a_2}, [..., [T^{a_{n-1}}, T^{a_n}]...]]). \tag{1.13}$$

With the imposed relations (1.12) and that partial amplitudes $\widetilde{A}(1, \sigma, n)$ are cyclic and satisfy KK-relation, we see that the claim is true.

Above logic is perfectly right, but there are several unclear points. The first thing is that each solution is based on (1.12) where two numbers have been fixed. Choosing different pairs, we will have different solutions in principle. For example, the same $\tau_{123..n}$-function can have two different expressions with two different choices of fixed pair (i_1, j_1) and (i_2, j_2). Secondly, for solutions with fixed pair, for example, $(1, n)$, does two τ-functions have **natural relabeling property**? More explicitly, assuming two τ-functions have expressions $\tau_{\sigma_1} = \sum_i c_i n_{\alpha_i}$ and $\tau_{\sigma_2} = \sum_j d_j n_{\beta_j}$, if $\sigma_2 = P(\sigma_1)$, where P is a permutation of n-elements, do we have

$$\sum_j d_j n_{\beta_j} \overset{?}{=} \sum_i c_i n_{P(\alpha_i)} \ . \tag{1.14}$$

There two points are related to each other. In fact, if (1.14) is satisfied, it should be expected that τ-functions are unique no matter which pair has been fixed in (1.12) to get them, since different choices can be related to each other by a permutation. In this paper, we will show that solutions obtained by our algorithm based on (1.12) will have the natural relabeling property. In other words, these two different construction methods are in fact consistent with each other. With this understanding, we show that how to use the relabeling property to fix τ-functions.

The structure of this paper is the following. In section 2, we provide a short review of the kinematic algebra proposed in [29]. Then we provide an algorithm to construct dual-trace factors in section 3. To demonstrate the idea outlined in section 3, we present several examples in section 4. We discuss the natural relabeling property and prove that the solutions obtained from our algorithm do satisfy the relabeling symmetry. Finally, a summary of this work is given in section 6.



Figure 1: The Jacobi identity for kinematic structure constant. It is given by sum over cyclic ordering of three incoming arrows $1, 2, 3$.

2. Useful properties of BCJ numerators

Before presenting the construction of dual-trace factors, we review some useful properties of BCJ numerator discussed in [29].

The generator of kinematic algebra is given by

$$T^{k,a} \equiv e^{ik\cdot x}\partial_a, \tag{2.1}$$

and the commutation relation gives the kinematic structure constant f

$$\begin{aligned}[T^{k_1,a}, T^{k_2,b}] &= (-i)(\delta_a{}^c k_{1b} - \delta_b{}^c k_{2a})\, e^{i(k_1+k_2)\cdot x}\partial_c \\ &= f^{(k_1,a),(k_2,b)}{}_{(k_1+k_2,c)}\, T^{(k_1+k_2,c)}.\end{aligned} \tag{2.2}$$

The kinematic structure constant satisfies antisymmetry property

$$f^{12}{}_3 = -f^{21}{}_3, \tag{2.3}$$

and Jacobi identity

$$f^{1_a,2_b}{}_{(1+2)^e} f^{(1+2)_e,3_c}{}_{(1+2+3)^d} + f^{2_b,3_c}{}_{(2+3)^e} f^{(2+3)_e,1_a}{}_{(1+2+3)^d} + f^{3_c,1_a}{}_{(1+3)^e} f^{(1+3)_e,2_b}{}_{(1+2+3)^d} = 0. \tag{2.4}$$

It is worth to notice that unlike the structure constant of gauge group, for $f^{12}{}_3$ here there is no natural way to put index 3 into same footing as index $1, 2$, thus to distinguish these three indices, we can use arrows and the Jacobi identity can be drawn in Figure 1 as the cyclic sum over incoming three arrows.

From our previous work [29], the total amplitude can be written as dual-DDM form where the **nu-**

merator factor is given as[1]

$$n_{12\ldots(n-1)n}=\sum_{j=1}^{N}c_j\epsilon(q_j)\cdot$$



(2.5)

where each term in the bracket is constructed by kinematic structure constant. Because the Jacobi identity (2.4) and antisymmetry property (2.3), the numerator factor n will have some very nice properties. For example, we will have following two identities

$$\begin{aligned}&n_{n\alpha_1\ldots\alpha_i1\rho_1\ldots\rho_j(n-1)}\\ =&\sum_{\textit{splittings }O\{\rho_1,\ldots,\rho_j\}\to O\{\beta_1,\ldots,\beta_r\},O\{\gamma_1,\ldots,\gamma_s\}}(-1)^{r+1}n_{n\alpha_1\ldots\alpha_i\gamma_1\ldots\gamma_s(n-1)\beta_r\ldots\beta_11}.\end{aligned}\tag{2.6}$$

and

$$\begin{aligned}&n_{n\alpha_1\ldots\alpha_i1\rho_1\ldots\rho_j(n-1)}\\ =&\sum_{\textit{splittings }O\{\rho_1,\ldots,\rho_j\}\to O\{\beta_1,\ldots,\beta_r\},O\{\gamma_1,\ldots,\gamma_s\}}(-1)^{i+s}n_{1\beta_1\ldots\beta_r(n-1)\gamma_s\ldots\gamma_1\alpha_i\ldots\alpha_1n}.\end{aligned}\tag{2.7}$$

where the sum in (2.6) and (2.7) is over all possible splitting of the set ρ into two subsets β and γ (both sets β and γ can be empty set). In each subset, the relative ordering must be same as the relative ordering in mother set ρ. These two identities are relabeling invariant, i.e., if we act a permutation $P\in S_n$ on sub-indices at both sides, there two identities still hold.

These two identities are very important to show the relabeling property of τ-functions constructed by the algorithm in previous section. Thus we want to give a proof. Since the $c_j\epsilon(q_j)$-part is same, we just need to prove the part in the bracket of (2.5) satisfies above two identities. For this purpose we draw the typical term in part (a) of Figure 2 and apply the Jacobi identity to the part indicated by square. The

---

[1] Since in our whole paper, we will use numerator factor of the DDM-chain form only, to simply the notation we will use $n_{123\ldots n}$.



(a) A typical term



(b) Applying Jacobi relation for particular arrows

Figure 2: (a) A typical term for $n_{n\alpha_1\ldots\alpha_i 1\rho_1\ldots\rho_j(n-1)}$ where we have used square to indicate the place to apply Jacobi identity. (b) The manipulation using Jacobi identity with particular arrow assignment.

result is give in part (b) of Figure 2 for a particular arrow configuration. It can be shown that same result will be obtained for all other possible arrow configurations. The result in part (b) tells us that we can move the ρ_1 attached to the $(n-1)$-th block to two places: the first place is attached to n-th block with $+$ sign and the second place is attached to 1 with $-$ sign. Repeating above manipulations, we can move down $\rho_1, \rho_2, ..., \rho_j$. The final configuration will be the ordering $\{n, \{\alpha_1, ..., \alpha_i\}, \{\gamma_1, ..., \gamma_s\}, n-1, \{\beta_r, ..., \beta_1\}, 1\}$ with sign $(-)^{r+1}$ (The extra $-$ sign comes from pulling $n-1$ from up to down by antisymmetry), where sets $\{\gamma_1, ..., \gamma_s\}$ and $\{\beta_1, ..., \beta_r\}$ is the splitting of original set $\{\rho_1, ..., \rho_j\}$ with relative ordering kept in each subset. By now, we have proved the identity (2.6). Finally we reverse the ordering to get $\{1, \{\beta_1, ..., \beta_r\}, n-1, \{\gamma_s, ..., \gamma_1\}, \{\alpha_i, ..., \alpha_1\}, n\}$ with sign $(-)^{r+1+(n-2)}$. Using $n-3 = i+j$ and $j = r+s$ we get the sign to be $(-)^{i+s}$. Thus the identity (2.7) is proved.

It can be shown that identity (2.6) and (2.7) can be replaced by following two forms (where for later application we have switched the $1 \leftrightarrow n$)

$$n_{1\alpha_1\ldots\alpha_i n\rho_1\ldots\rho_j(n-1)} = -n_{1\alpha_1\ldots\alpha_i[\rho_1,[\rho_2,\ldots[\rho_j,(n-1)]\ldots]]n}. \tag{2.8}$$

and

$$n_{\rho_1\ldots\rho_j 1,\alpha_1,\ldots,\alpha_i,n} = (-)\, n_{1[[[\rho_1,\rho_2],\ldots],\rho_j],\alpha_1,\alpha_i,n}\,, \tag{2.9}$$

where [,] is again the anti-commutative bracket. If the set ρ is empty, $[\rho_1,[\rho_2,...[\rho_j,(n-1)]...]]$ is just $(n-1)$. Formula (2.8) and (2.9) can be used to move one index to the last or to the first position.

Before we end this section, there is one remark. It will be clear soon that all discussions in this paper will based on above two identities (2.6) and (2.7). We have proved it using the explicit construction (2.5), where the local diagram picture is manifest. However, since our proof relies only the Jacobi identity of BCJ numerator n, it seems that our local diagram picture is not necessary and the result will be true in general by conditions imposed on n.

3. Construction of τ by BCJ numerators

In this section, we present the construction of τ-functions using (1.12), (1.10) and (1.11). With $(1,n)$ fixed and a given permutation σ_2,...,σ_{n-2}, the relation between τ and n is given by (1.12)

$$n_{1\sigma_2\ldots\sigma_{n-1}n} = \tau_{1[\sigma_2,[\ldots,[\sigma_{n-1},n]\ldots]]}, \tag{3.1}$$

where the bracket $[A,B]\equiv AB-BA$. For example, we have

$$\tau_{1[2,[3,4]]} = \tau_{1234} - \tau_{1243} - \tau_{1342} + \tau_{1432}.$$

The expansion at the right handed side of (3.1) includes cases where n is not at the last position. Thus we should use KK-relation (1.11) to reduce them. After such reduction we will get

$$n_{1\sigma_2\ldots\sigma_{n-1}n} = \sum_{\sigma'\in S_{n-2}} G_{1,n}(\sigma|\sigma')\tau_{1\sigma' n}. \tag{3.2}$$

To solve τ by n, we need to understand the matrix $G_{1,n}(\sigma|\sigma')$. The $(n-2)!\times(n-2)!$ matrix can be obtained by following manipulations. First it is easy to see that

$$\tau_{1[\sigma_2,[\ldots,[\sigma_{n-1},n]\ldots]]} = \sum_{splittings\ \sigma\to\alpha,\beta} (-1)^{n_\beta}\tau_{1\alpha n\beta^T} \tag{3.3}$$

where the sum is over all possible splitting of ordered set $\{\sigma_2,...,\sigma_{n-1}\}$ into two subsets α and β (empty set is allowed) such that inside each subset the relative ordering inside the set σ has been kept. For example, the set $\sigma=234$ has following eight splittings

$$(\alpha,\beta) = (\{234\},\emptyset)/(\{23\},\{4\})/(\{24\},\{3\})/(\{34\},\{2\})/(\emptyset,\{234\})/(\{4\},\{23\})/(\{3\},\{24\})/(\{2\},\{34\})$$

Secondly using the imposed KK relation on τ, any $(-1)^{n_\beta}\tau_{1\alpha n\beta^T}$ can be expressed as $\sum\limits_{\rho\in OP(\{\alpha\}\bigcup\{\beta\})}\tau_{1\rho n}$. Combining these two steps we arrive

$$\tau_{1[\sigma_2,[\ldots,[\sigma_{n-1},n]\ldots]]} = \sum_{splittings\ \sigma\to\alpha,\beta}\ \sum_{\rho\in OP(\{\alpha\}\bigcup\{\beta\})}\tau_{1\rho n}. \tag{3.4}$$

From this formula we can see that that $G_{1,n}(\sigma|\sigma')$ is the number of splittings of σ into two subsets α, β, such that ordering σ' can be obtained by recombining two subsets α, β arbitrarely with relative ordering kept inside each subset.

Let us give a few examples of $G_{1,n}(\sigma|\sigma')$ with $\sigma = \{234\}$. For $\sigma' = \{234\}$, there are eight splitting of σ, which can be used to recombine to get σ', so $G_{1,5}(\{234\}|\{234\}) = 8$. For $\sigma' = \{243\}$ there are four splittings $(\{23\},\{4\})$, $(\{24\},\{3\})$, $(\{4\},\{23\})$, $(\{3\},\{24\})$, so $G_{1,5}(\{234\}|\{243\}) = 4$. For $\sigma' = \{423\}$ there are only two splittings $(\{23\},\{4\})$, $(\{4\},\{23\})$, so $G_{1,5}(\{234\}|\{423\}) = 2$. For $\sigma' = \{432\}$, there is no any splitting, so $G_{1,5}(\{234\}|\{432\}) = 0$. In fact, one can show that for three numbers i, j, k, if their ordering inside the set σ is $i > j > k$ and their ordering inside the set σ' is $i < j < k$, the $G_{1,n}(\sigma|\sigma') = 0$. The reason is following. To reproduce the right ordering inside σ', when we split σ into two subsets, pair i, j can not belong to same subsets. Same conclusions hold for pairs i, k and j, k, which can not be satisfied. With this general arguments, we can see that majority elements of matrix $G_{1,n}(\sigma|\sigma')$ are zero.

Now we discuss some general properties of matrix $G_{1,n}(\sigma|\sigma')$:

- (1) First the value can be written as

  $$G_{1,n}(\sigma|\sigma') = \sum_{s' \in \{splittings\ \sigma' \to \alpha,\beta'\}} \sum_{s \in \{splittings\ \sigma \to \alpha,\beta\}} \delta(s|s'), \tag{3.5}$$

  where s and s' are the splittings of σ and σ' into two ordered sets respectively. This means elements of matrix $G_{1,n}(\sigma|\sigma')$ can be calculated by another way. First we find all 2^{n_σ} splittings of the set σ into two subsets (α, β) with relative ordering kept and all splittings of the set σ' into two subsets (α', β') with relative ordering kept. Then we find all subsets such that $(\alpha, \beta) = (\alpha', \beta')$. The total number is $G_{1,n}(\sigma|\sigma')$.

- (2) From the above property (A.11), it is easy to see that G is symmetric

  $$G_{1,n}(\sigma|\sigma') = G_{1,n}(\sigma'|\sigma). \tag{3.6}$$

- (3) There are some special values. When $\sigma = \sigma'$, all possible splittings of $\sigma \in S_{n-2}$ are included, i.e., $G_{1,n}(\sigma|\sigma) = 2^{n-2}$. If σ, σ' are different only by one permutation of two adjacent numbers, $G_{1,n}(\sigma|\sigma) = 2^{n-3}$. If $\sigma = (..., i_1, i_2, ..., j_1, j_2, ...)$ and $\sigma' = (..., i_2, i_1, ..., j_2, j_1, ...)$, $G_{1,n}(\sigma|\sigma) = 2^{n-4}$. Similar pattern holds for more pairs interchangings.

- (4) The fourth observation is that

  $$G_{1,n}(\sigma|\sigma') = G_{1,n}(P(\sigma)|P(\sigma')) \tag{3.7}$$

  where P is any permutation. The reason is that $G_{1,n}(\sigma|\sigma')$ care only the relative positions between sets σ, σ', so it does not matter if a element is called x or y.

Having discussed the matrix $G_{1,n}(\sigma|\sigma')$, we can solve $\tau_{1,\sigma,n}$ by BCJ numerator factors using (3.2) and obtain

$$\tau_{1\sigma' n}=\frac{\det\mathbb{G}'}{\det\mathbb{G}}, \tag{3.8}$$

where $\mathbb{G}$ is an $(n-2)!\times(n-2)!$ matrix with elements $G_{1,n}(\sigma|\sigma')$, and $\mathbb{G}'$ is obtained by replacing the column σ' of $\mathbb{G}$ by the column of BCJ numerator factor $n_{1\sigma n}$. Now we have provide a general construction of the $(n-2)!$ τ's with 1, n fixed at the two ends (3.8). Other $(n!-(n-2)!)$ τ's can be obtained by KK relation and cyclic symmetry (see (1.10) and (1.11)).

Our above construction for τ's of arbitrary tree amplitudes is definite and there is no any ambiguity. Only subtle point is that in above construction, two special elements have been fixed, for example 1 and n. Then it is natural to ask what is the relation between two solutions obtained by fixed two different pairs of elements (i_1,j_1) and (i_2,j_2)? There are two possibilities: one is that these two solutions do not have any relation and another is that these two solutions will relate to each other by some manipulations. As we will show in later sections, there is **natural relabeling property** for solution obtained by above construction. This property tells us that if we have an expression for one τ-function, we will get others by relabeling. Furthermore, all solutions with different fixed pairs will give same answer.

4. Examples

In this section, we will use several examples to demonstrate explicitly the algorithm for finding τ-functions and check that the solution satisfies the **natural relabeling property**.

4.1 Three-point case

In this case, with $1,3$ fixed we have $G_{1,3}(\{2\}|\{2\})=2$,

$$n_{123}=2\tau_{123}\Longrightarrow\tau_{123}=\frac{1}{2}n_{123}. \tag{4.1}$$

There are other five τ's can be obtained from above by cyclic symmetry and KK-relation

$$\begin{aligned} Cyclic: &\quad \tau_{312}=\tau_{231}=\tau_{123}\\ KK-rel: &\quad \tau_{213}=\tau_{321}=\tau_{132}=-\tau_{123} \end{aligned} \tag{4.2}$$

To check the relabeling, noticing that if we exchange 3 ↔ 2 in (4.1), we will get

$$\widetilde{\tau}_{132}=\frac{1}{2}n_{132} \tag{4.3}$$

Because for three point case, BCJ numerator n has cyclic symmetry as well as $n_{123}=-n_{132}$ by construction, we see that $\widetilde{\tau}_{132}$ is equal to τ_{132} in (4.2), i.e., it does have the relabeling property. It is easy to check the relabeling property for other τ's.

4.2 Four-point case

For four point case, we have $4! = 24$ τ's. Now we use our algorithm to determine all of them and check the relabeling property!

Solving the τ's in KK-basis: we have $(4-2)! = 2$ equations and $(4-2)! = 2$ independent τ's. With ordering $(\{23\},\{32\})$ the matrix is

$$G_{1,4} = \begin{pmatrix} 4 & 2 \\ 2 & 4 \end{pmatrix} \tag{4.4}$$

From this we can solve

$$\tau_{1234} = \frac{\begin{vmatrix} n_{1234} & 2 \\ n_{1324} & 4 \end{vmatrix}}{\begin{vmatrix} 4 & 2 \\ 2 & 4 \end{vmatrix}} = \frac{1}{3}n_{1234} - \frac{1}{6}n_{1324} \tag{4.5}$$

and

$$\tau_{1324} = \frac{\begin{vmatrix} 4 & n_{1234} \\ 2 & n_{1324} \end{vmatrix}}{\begin{vmatrix} 4 & 2 \\ 2 & 4 \end{vmatrix}} = -\frac{1}{6}n_{1234} + \frac{1}{3}n_{1324} \tag{4.6}$$

Find remaining τ's: Having knowing τ_{1234} and τ_{1324}, we can construct other 22 τ's. Using imposed KK-relation, we can obtain following four τ with 1 fixed:

$$\begin{aligned}
\tau_{1243} &\equiv -\tau_{1234} - \tau_{1324} = -\frac{1}{6}n_{1234} - \frac{1}{6}n_{1324} \\
\tau_{1342} &\equiv -\tau_{1234} - \tau_{1324} = -\frac{1}{6}n_{1234} - \frac{1}{6}n_{1324} \\
\tau_{1423} &\equiv +\tau_{1324} = -\frac{1}{6}n_{1234} + \frac{1}{3}n_{1324} \\
\tau_{1432} &\equiv +\tau_{1234} = \frac{1}{3}n_{1234} - \frac{1}{6}n_{1324}
\end{aligned} \tag{4.7}$$

Having obtained all $\tau_{1\sigma}$, the remaining 18 τ's are obtained by imposed cyclic symmetry. For example, we have

$$\tau_{1234} = \tau_{4123} = \tau_{3412} = \tau_{2341} = \frac{1}{3}n_{1234} - \frac{1}{6}n_{1324} \tag{4.8}$$

Relabeling property: Within explicit result, we can check the relabeling property. First it is easy to see that expression (4.6) can be obtained from (4.5) by replacing 2 → 3, 3 → 2. In principal, this fact does not need to hold. However, by symmetric property and and property (3.7) of matrix G, it is natural to get $\tau_{1P(\sigma)n} = P(\tau_{1\sigma n})$.

Now we check the relabeling property for $\tau_{1\sigma}$. From solution τ_{1234} by index relabeling $(3,4) \to (4,3)$ we can write down

$$\widetilde{\tau}_{1243} = \frac{1}{3} n_{1243} - \frac{1}{6} n_{1423} \tag{4.9}$$

To check if $\widetilde{\tau}_{1243}$ is equal to τ_{1243} obtained in (4.7), we use the relation (2.6) to get

$$n_{1243} = -n_{1234}, \quad n_{1423} = -n_{1234} + n_{1324} \tag{4.10}$$

Putting it back, it is easy to check $\widetilde{\tau}_{1243} = \tau_{1243}$. In other words, two expressions, i.e., the one obtained by our imposed KK-relation and the one obtained by index relabeling from known solution $\tau_{1\sigma 4}$ are same!

Next, we check the relabeling property for τ's where 1 is not the first element. By relabeling $(1,2,3,4) \to (4,1,2,3)$ from τ_{1234} we have

$$\widetilde{\tau}_{4123} = \frac{1}{3} n_{4123} - \frac{1}{6} n_{4213}. \tag{4.11}$$

Using the relation (2.7) to expand

$$n_{4123} = n_{1234} - n_{13(24)}, \quad n_{4213} = -n_{1324}, \tag{4.12}$$

we do find $\widetilde{\tau}_{4123}$ is equal to τ_{4123} written down in (4.8).

Although we have present only three examples, using the Mathematica we do check that indeed result obtained from our algorithm (including using KK and cyclic relations) does agree with the one obtained from relabeling of single τ_{1234} solution.

4.3 five-point case

Having above four point example, we will not give too many details for five, six and seven points.

Solutions from our algorithm: First we can solve τ's in KK-basis using following equation

$$\begin{pmatrix} n_{12345} \\ n_{12435} \\ n_{13245} \\ n_{14235} \\ n_{13425} \\ n_{14325} \end{pmatrix} = \begin{pmatrix} 2^3 & 2^2 & 2^2 & 2^1 & 2^1 & 0 \\ 2^2 & 2^3 & 2^1 & 2^2 & 0 & 2^1 \\ 2^2 & 2^1 & 2^3 & 0 & 2^2 & 2^1 \\ 2^1 & 2^2 & 0 & 2^3 & 2^1 & 2^2 \\ 2^1 & 0 & 2^2 & 2^1 & 2^3 & 2^2 \\ 0 & 2^1 & 2^1 & 2^2 & 2^2 & 2^3 \end{pmatrix} \begin{pmatrix} \tau_{12345} \\ \tau_{12435} \\ \tau_{13245} \\ \tau_{14235} \\ \tau_{13425} \\ \tau_{14325} \end{pmatrix}. \tag{4.13}$$

and obtain the expression τ_{12345} by BCJ numerator factor as

$$\tau_{12345} = \frac{1}{4} n_{12345} - \frac{1}{10} n_{12435} - \frac{1}{20} n_{14235} - \frac{1}{10} n_{13245} - \frac{1}{20} n_{13425} + \frac{1}{10} n_{14325}. \tag{4.14}$$

Other five τ's can be obtained by relabeling $2, 3, 4$ from (4.14). Having these six τ solutions for KK-basis, we can use the KK-relation and cyclic relation to find other 114 τ's.

Relabeling property: Now we need to check if above result has the relabeling property. Although we have used Mathematica to check that all 120 τ's are related to each other by relabeling property, here we will give only a few examples. The first example is[2]

$$\tau_{12534} = -\frac{1}{20}n_{12345} + \frac{1}{10}n_{12435} - \frac{1}{20}n_{13425} + \frac{1}{20}n_{14235} + \frac{1}{10}n_{14325} \tag{4.15}$$

which is obtained from KK-relation. Now we relabel the solution (4.14) with $3 \to 5, 4 \to 3, 5 \to 4$. After using

$$\begin{aligned}
n_{12534} &= -n_{12345} + n_{12435} \\
n_{12354} &= -n_{12345} \\
n_{13254} &= -n_{13245} \\
n_{15234} &= -n_{12345} + n_{12435} + n_{13425} - n_{14325} \\
n_{15324} &= n_{12435} - n_{13245} + n_{13425} - n_{14235} \\
n_{13524} &= -n_{13245} + n_{13425}
\end{aligned}$$

it can be checked that the relabeling (4.14) does reproduce the (4.15).

The second example is $\tau_{51234} \equiv \tau_{12345}$ by our definition. The relabeling of (4.14) gives

$$\widetilde{\tau}_{51234} = \frac{1}{4}n_{51234} - \frac{1}{10}n_{51324} - \frac{1}{20}n_{53124} - \frac{1}{10}n_{52134} - \frac{1}{20}n_{52314} + \frac{1}{10}n_{53214}. \tag{4.16}$$

Using the result

$$\begin{aligned}
n_{51234} &= n_{12345} - n_{12435} - n_{13425} + n_{14325}, \\
n_{51324} &= n_{13245} - n_{13425} - n_{12435} + n_{14235}, \\
n_{53124} &= -n_{12435} + n_{14235}, \\
n_{52134} &= -n_{13425} + n_{14325}, \\
n_{52314} &= n_{14325}, \\
n_{53214} &= n_{14235}.
\end{aligned} \tag{4.17}$$

it is easy to see that $\tau_{51234} = \widetilde{\tau}_{51234}$, i.e., the relabeling property is satisfied.

---

[2] It is worth to notice that the expression of τ_{12534} is simpler than (4.14) because it has only 5 terms with simpler coefficients. It will be interesting to investigate if it is a general thing.

4.4 Six-point case

For six points, using our algorithm we can solve 24 τ's in the KK-basis. Since they are related to each other by relabeling, we can write down only one solution τ_{123456}

$$\begin{aligned}\tau_{123456} = \frac{1}{630}\Big[&126n_{123456} - 44n_{123546} - 44n_{124356} - 19n_{124536} - 19n_{125346} + 36n_{125436}\\ &-44n_{132456} + 16n_{132546} - 19n_{134256} - 19n_{134526} + n_{135246} + 11n_{135426}\\ &-19n_{142356} + n_{142536} + 36n_{143256} + 11n_{143526} - 9n_{145236} + 16n_{145326}\\ &-19n_{152346} + 11n_{152436} + 11n_{153246} + 16n_{153426} + 16n_{154236} - 44n_{154326}\Big].\end{aligned}\tag{4.18}$$

Using the KK-basis of τ we can get remaining 696 τ's. Now we check the relabeling property. We have used the Mathematica to check that all 720 τ's are related to each other by relabeling property. Here we give only one example $\tau_{612345} \equiv \tau_{123456}$ by cyclic symmetry. The relabeling from τ_{123456} gives

$$\begin{aligned}\widetilde{\tau}_{612345} = \frac{1}{630}\Big[&126n_{612345} - 44n_{612435} - 44n_{613245} - 19n_{613425} - 19n_{614235} + 36n_{614325}\\ &-44n_{621345} + 16n_{621435))} - 19n_{623145} - 19n_{623415} + n_{624135} + 11n_{624315}\\ &-19n_{631245} + n_{631425} + 36n_{632145} + 11n_{632415} - 9n_{634125} + 16n_{634215}\\ &-19n_{641235} + 11n_{641325} + 11n_{642135} + 16n_{642315} + 16n_{643125} - 44n_{643215}\Big].\end{aligned}\tag{4.19}$$

Using following result

$$\begin{aligned}n_{61ijk5} &= n_{1ijk56} - n_{1jk5i6} - n_{1ik5j6} - n_{1ij5k6} + n_{1k5ji6} + n_{1i5kj6} + n_{1j5ki6} - n_{15kji6}\\ n_{6i1jk5} &= -n_{1jk5i6} + n_{1j5ki6} + n_{1k5ji6} - n_{15kji6}\\ n_{6ij1k5} &= n_{1k5ji6} - n_{15kji6}\\ n_{6ijk15} &= -n_{15kji6}.\end{aligned}\tag{4.20}$$

We can see $\widetilde{\tau}_{612345} = \tau_{612345}$.

4.5 Seven-point case

For seven-points, the expression $\tau_{1234567}$ with $1, 7$ fixed will be

$$\begin{aligned}\tau_{1234567} = \tfrac{1}{24192}\Big(&4032n_{1234567} - 1284n_{1234657} - 1284n_{1235467} - 513n_{1235647} - 513n_{1236457} + 940n_{1236547}\\ &-1284n_{1243567} + 393n_{1243657} - 513n_{1245367} - 438n_{1245637} + 27n_{1246357} + 259n_{1246537}\\ &-513n_{1253467} + 27n_{1253647} + 940n_{1254367} + 259n_{1254637} - 213n_{1256347} + 337n_{1256437}\\ &-438n_{1263457} + 259n_{1263547} + 259n_{1264357} + 344n_{1264537} + 337n_{1265347} - 940n_{1265437}\\ &-1284n_{1324567} + 421n_{1324657} + 393n_{1325467} + 205n_{1325647} + 205n_{1326457} - 337n_{1326547}\end{aligned}$$

$$
\begin{aligned}
&-513n_{1342567}+205n_{1342657}-438n_{1345267}-513n_{1345627}+32n_{1346257}+205n_{1346527}\\
&+27n_{1352467}-56n_{1352647}+259n_{1354267}+184n_{1354627}+23n_{1356247}+157n_{1356427}\\
&+32n_{1362457}-6n_{1362547}-51n_{1364257}+143n_{1364527}+6n_{1365247}-259n_{1365427}\\
&-513n_{1423567}+205n_{1423657}+27n_{1425367}+32n_{1425637}-56n_{1426357}-6n_{1426537}\\
&+940n_{1432567}-337n_{1432657}+259n_{1435267}+205n_{1435627}-6n_{1436257}-148n_{1436527}\\
&-213n_{1452367}+23n_{1452637}+337n_{1453267}+157n_{1453627}-213n_{1456237}+337n_{1456327}\\
&+23n_{1462357}-53n_{1462537}+6n_{1463257}+11n_{1463527}+148n_{1465237}-205n_{1465327}\\
&-438n_{1523467}+32n_{1523647}+259n_{1524367}-51n_{1524637}+23n_{1526347}+6n_{1526437}\\
&+259n_{1532467}-6n_{1532647}+344n_{1534267}+143n_{1534627}-53n_{1536247}+11n_{1536427}\\
&+337n_{1542367}+6n_{1542637}-940n_{1543267}-259n_{1543627}+148n_{1546237}-205n_{1546327}\\
&-213n_{1562347}+148n_{1562437}+148n_{1563247}+107n_{1563427}+107n_{1564237}-393n_{1564327}\\
&-513n_{1623457}+205n_{1623547}+184n_{1624357}+143n_{1624537}+157n_{1625347}-259n_{1625437}\\
&+205n_{1632457}-148n_{1632547}+143n_{1634257}+344n_{1634527}+11n_{1635247}-184n_{1635427}\\
&+157n_{1642357}+11n_{1642537}-259n_{1643257}-184n_{1643527}+107n_{1645237}-421n_{1645327}\\
&+337n_{1652347}-205n_{1652437}-205n_{1653247}-421n_{1653427}-393n_{1654237}+1284n_{1654327}\Big).
\end{aligned}
\tag{4.21}
$$

By relabeling the above expression $(1,2,3,4,5,6,7) \to (7,1,2,3,4,5,6)$, we get $\widetilde{\tau}_{7123456}$. To show that it is equal to $\tau_{7123456} = \tau_{1234567}$ from cyclic symmetry, we just need to use following expressions

$$
\begin{aligned}
n_{71ijkl6} &= n_{1ijkl67}-n_{1ijk6l7}-n_{1ijl6k7}+n_{1ij6lk7}-n_{1ikl6j7}+n_{1ik6lj7}+n_{1il6kj7}-n_{1i6lkj7}\\
&\quad - n_{1jkl6i7}+n_{1jk6li7}+n_{1jl6ki7}-n_{1j6lki7}+n_{1kl6ji7}-n_{1k6lji7}-n_{1l6kji7}+n_{16lkji7}\\
n_{7i1jkl6} &= -n_{1jkl6i7}+n_{1jk6li7}+n_{1jl6ki7}-n_{1j6lki7}+n_{1kl6ji7}-n_{1k6lji7}-n_{1l6kji7}+n_{16lkji7}\\
n_{7ij1kl6} &= n_{1kl6ji7}-n_{1k6lji7}-n_{1l6kji7}+n_{16lkji7}\\
n_{7ijk1l6} &= -n_{1l6kji7}+n_{16lkji7}\\
n_{7ijkl16} &= n_{16lkji7}
\end{aligned}
\tag{4.22}
$$

5. Relabeling property

In previous section, we have demonstrated the consistency between our algorithm and the relabeling property. In this section, we will give a general understanding of this property. Before doing so, we need to address a technical issue, i.e., the definition of matrix G. Since the relabeling is a permutation of n

elements while the definition of G involves only permutation of $(n-2)$ elements after two elements having been fixed. To relate matrix G with different fixed pairs, we enlarge the definition of matrix $G_{1,n}[\sigma|\sigma']$ to $G[\{1,\sigma,n\}|1,\sigma',n]$ and require that when we split the set $\{1,\sigma,n\}$ to two subsets, the first element 1 must be at the first subset and the last element n must be at the second subset. It is easy to see that the new definition is equivalent to the old one. More importantly, the relabeling property (3.7) with $(n-2)$ elements can be enlarged to relabeling property with n elements, i.e, we will have

$$G[\sigma|\sigma'] = G[P(\sigma)|P(\sigma')], \qquad P \in S_n \; , \tag{5.1}$$

where σ, σ' are lists with n elements.

Having enlarged the definition of matrix G, we can start our discussion. The structure of this section is the following. First we will set up the general framework of discussion. Then we will discuss how the relabeling property can be used to solve τ-function.

5.1 The proof of relabeling invariance

We want to prove the relabeling property without knowing the solution explicitly. Since by our algorithm three equations (1.12), (1.10) and (1.11) fix the solution uniquely, we can prove the relabeling property by showing that above three equations have relabeling property. The imposed cyclic symmetry (1.10) and KK-relation (1.11) have the property obviously, thus the key is to show that equation (1.12) does have the relabeling property.

Now we consider the relabeling $x_s \to z_s$ with $s = 1, ..., n$. Under the relabeling, an ordering of n elements becomes another ordering of n elements, i.e., $X_i \to Z_i$ and equation (1.12) is relabeled to

$$\sum_j G_{X_i X_j} \tau_{X_j} = n_{X_i} \Longrightarrow \sum_j G_{Z_i Z_j} \tau_{Z_j} = n_{Z_i} \tag{5.2}$$

However, by our cyclic and KK-relation, we have

$$\tau_{Z_i} = Q_{Z_i X_i} \tau_{X_i} \tag{5.3}$$

and by (2.7) and (2.6) we have

$$n_{Z_i} = P_{Z_i X_i} n_{X_i} \tag{5.4}$$

thus

$$G_{Z_i Z_j} \tau_{Z_j} = n_{Z_i} \Longrightarrow G_{Z_i Z_j} Q_{Z_i X_i} \tau_{X_i} = P_{Z_i X_i} n_{X_i} \tag{5.5}$$

To ensure solution from our algorithm to have relabeling property, we just need to have

$$[P_{Z_i X_i}]^{-1} G_{Z_i Z_j} Q_{Z_j X_j} = G_{X_i X_j} \tag{5.6}$$

Now let us demonstrate the use of (5.6) by several examples. First let us consider the four point case where the matrix G is given by (4.4). For relabeling given by permutation $P(3,4)$, we have $X_1 = (1,2,3,4) \to Z_1 = (1,2,4,3)$ and $X_2 = (1,3,2,4) \to Z_2 = (1,4,2,3)$. Under this permutation we can find

$$\begin{pmatrix} \tau_{Z_1} \\ \tau_{Z_2} \end{pmatrix} = \begin{pmatrix} -1 & -1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \tau_{X_1} \\ \tau_{X_2} \end{pmatrix} \tag{5.7}$$

and

$$\begin{pmatrix} n_{Z_1} \\ n_{Z_2} \end{pmatrix} = \begin{pmatrix} -1 & 0 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} n_{X_1} \\ n_{X_2} \end{pmatrix} \tag{5.8}$$

Thus it is easy to check that

$$\begin{pmatrix} 4 & 2 \\ 2 & 4 \end{pmatrix} = \begin{pmatrix} -1 & 0 \\ -1 & 1 \end{pmatrix}^{-1} \begin{pmatrix} 4 & 2 \\ 2 & 4 \end{pmatrix} \begin{pmatrix} -1 & -1 \\ 0 & 1 \end{pmatrix} \tag{5.9}$$

For the second example, we take permutation $P(1,2)$ for five point case and the G matrix is given by (4.13). With this permutation, we find the matrix P by following manipulation

$$\begin{pmatrix} n_{12345} \\ n_{12435} \\ n_{13245} \\ n_{13425} \\ n_{14235} \\ n_{14325} \end{pmatrix} \to \begin{pmatrix} n_{21345} \\ n_{21435} \\ n_{23145} \\ n_{23415} \\ n_{24135} \\ n_{24315} \end{pmatrix} = (-) \begin{pmatrix} n_{12345} \\ n_{12435} \\ n_{1[2,3]45} \\ n_{1[[2,3],4]5} \\ n_{1[2,4]35} \\ n_{1[[2,4],3]5} \end{pmatrix} = \begin{pmatrix} -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 \\ -1 & 0 & 1 & 0 & 0 & 0 \\ -1 & 0 & 1 & 0 & 1 & -1 \\ 0 & -1 & 0 & 0 & 1 & 0 \\ 0 & -1 & 1 & -1 & 1 & 0 \end{pmatrix} \begin{pmatrix} n_{12345} \\ n_{12435} \\ n_{13245} \\ n_{13425} \\ n_{14235} \\ n_{14325} \end{pmatrix}. \tag{5.10}$$

Similar manipulation gives the matrix Q as

$$P_{Z_iX_i} = \begin{pmatrix} -1 & 0 & -1 & -1 & 0 & 0 \\ 0 & -1 & 0 & 0 & -1 & -1 \\ 0 & 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & -1 & 0 & 0 \end{pmatrix}. \tag{5.11}$$

It is straightforward to see that $[P_{Z_iX_i}]^{-1} G_{Z_iZ_j} Q_{Z_jX_j} = G_{X_iX_j}$ is indeed satisfied.

Having shown two examples, now we discuss how we could give a general proof. To do so, we need following observation. Assuming the relabeling from $X_i \to Z_i$ is done by two steps $X_i \to Y_i$ and $Y_i \to Z_i$, it is easy to see that

$$\tau_{Z_i} = Q_{Z_iY_i}\tau_{Y_i} = Q_{Z_iY_i}Q_{Y_iX_i}\tau_{X_i}, \quad n_{Z_i} = P_{Z_iY_i}n_{Y_i} = P_{Z_iY_i}P_{Y_iZ_i}n_{Z_i} \tag{5.12}$$

and the condition (5.6) becomes

$$[P_{Z_iY_i}P_{Y_iZ_i}]^{-1}G_{Z_iZ_j}Q_{Z_iY_i}Q_{Y_iX_i}=G_{X_iX_j}\quad . \tag{5.13}$$

If the relabeling property holds for two steps $X_i\to Y_i$ and $Y_i\to Z_i$, i.e.

$$[P_{Y_iX_i}]^{-1}G_{Y_iY_j}Q_{Y_jX_j}=G_{X_iX_j} \tag{5.14}$$

and

$$[P_{Z_iY_i}]^{-1}G_{Z_iZ_j}Q_{Z_jY_j}=G_{Y_iY_j} \tag{5.15}$$

then (5.13) is also hold. This observation reflects the structure of group. It means that if we can show (5.6) is true for all generators of permutation group, it will be true for whole group.

For permutation S_n, there are $(n-1)$ generators. For our convenience, we can choose following generators $(n-2)$ generators

$$P_n=(n1),\quad P_i=(i,i+1),\quad i=2,3,...,n-2; \tag{5.16}$$

plus any one permutation of the form $P_{1i}=(1,j)$ or $P_{ni}=(n,i)$ with $i=2,3,...,n-1$. For permutations P_i with $i=2,...,n-2$ they are permutations among KK-basis with $(1,n)$ fixed and it is easy to see that $P=Q$ and $P^{-1}=P^T$. By the general property (3.7), $P^{-1}GP=G[P(\sigma)|P(\sigma')]=G[\sigma|\sigma']$, i.e., the condition (5.6) is satisfied.

For permutation P_n, since $2,3...,n-2$ are invariant, we have

$$\tau_{n\sigma 1}=\tau_{1n\sigma}=(-)^{n-2}\tau_{1\sigma^T n},\quad n_{n\sigma 1}=(-)^{n-2}n_{1\sigma^T n} \tag{5.17}$$

and especially

$$G[\{1\sigma^T n\}|\{1\gamma^T n\}]=G[\{1\sigma n\}|\{1\gamma n\}] \tag{5.18}$$

thus the relabeling property for permutation P_n is proved. To finish the proof, we need to check the last permutation P_{1i} or P_{ni}. Since it is very complicated, we give the proof in the Appendix.

5.2 An application

Having shown that solution from our algorithm has the relabeling property, it is natural to ask if we can use the relabeling property to fix the solution. In this subsection, we will show that it can be done. There are two different approaches and we will discuss one by one.

For the first approach, we need to use the relabeling property and only one equation of the form (1.12). To demonstrate the idea, let us start with four point example. First we expand τ_{1234} into the basis $n_{1\sigma n}$ as

$$\tau_{1234}=\alpha n_{1234}+\beta n_{1324} \tag{5.19}$$

Using the relabeling property other τ's in the KK-basis will have similar expansion. For $n = 4$, thing is simple and we have only one

$$\tau_{1324} = \alpha n_{1324} + \beta n_{1234} \tag{5.20}$$

To completely fix α, β we need to use one relation

$$n_{1234} = 4\tau_{1234} + 2\tau_{1324} = (4\alpha + 2\beta)n_{1234} + (4\beta + 2\alpha)n_{1324} \tag{5.21}$$

Since each n is independent, we have following two equations

$$4\alpha + 2\beta = 1, \quad 4\beta + 2\alpha = 0 \Longrightarrow \beta = -\frac{1}{6}, \quad \alpha = \frac{1}{3} \tag{5.22}$$

For general n-points, we expand, for example, $\tau_{123\ldots(n-1)n}$ into the BCJ numerator basis $n_{1\sigma n}$ with $(n-2)!$ unknown variables α_i. Then we use the relabeling property to express all other τ's in the KK-basis using same set of variables α_i. Next we put it back to equation $n_{123\ldots n} = \tau_{1[2,[3,\ldots[n-1,n]]]}$ and read out $(n-2)!$ equations from coefficients of BCJ numerator basis $n_{1\sigma n}$. From these equations we can solve α_i.

Above first approach has not used the full potential of relabeling property. Now we present the second approach. To demonstrate, we use the five point case as an example. Expanding τ_{12345} into the BCJ basis we have

$$\tau_{12345} = \alpha_1 n_{12345} + \alpha_2 n_{12435} + \alpha_3 n_{13245} + \alpha_4 n_{13425} + \alpha_5 n_{14235} + \alpha_6 n_{14325} \tag{5.23}$$

Using the relabeling property we can write down the expansion of other five τ's in the KK-basis using same six variables α_i. Up to this step, it is same as in the first approach. To use then full potential, there are two relabelings $P(15)$ and $P(54)$ we need to discuss. For permutation $P(15)$, we have at one side

$$\tau_{12345} \to \tau_{52341} = \alpha_1 n_{52341} + \alpha_2 n_{52431} + \alpha_3 n_{53241} + \alpha_4 n_{53421} + \alpha_5 n_{54231} + \alpha_6 n_{54321} \tag{5.24}$$

by relabeling property and at the another side

$$\begin{aligned} \tau_{12345} \to \tau_{52341} &= \tau_{15234} = -\tau_{14325} \\ &= -(\alpha_1 n_{14325} + \alpha_2 n_{14235} + \alpha_3 n_{13425} + \alpha_4 n_{13245} + \alpha_5 n_{12435} + \alpha_6 n_{12345}) \end{aligned} \tag{5.25}$$

by the cyclic symmetry, KK-relation and the expansion of KK-basis. Comparing these two results using $n_{1\sigma n} = (-)^n n_{n\sigma^T 1}$, we get immediately following equations

$$\alpha_1 = \alpha_1, \quad \alpha_6 = \alpha_6, \quad \alpha_2 = \alpha_3, \quad \alpha_4 = \alpha_5 \tag{5.26}$$

In other words, using the permutation $P(15)$ we reduce six unknown variables to four unknown variables.

Now we discuss the permutation $P(45)$. At one side we have

$$\tau_{12345} \to \tau_{12354} = \alpha_1 n_{12354} + \alpha_2 n_{12534} + \alpha_3 n_{13254} + \alpha_4 n_{13524} + \alpha_5 n_{15234} + \alpha_6 n_{15324}$$

$$\begin{aligned}&= n_{12345}(-\alpha_1-\alpha_2-\alpha_5)+n_{12435}(\alpha_2+\alpha_5+\alpha_6)+n_{13245}(-\alpha_3-\alpha_4-\alpha_6)\\&\quad +n_{13425}(\alpha_4+\alpha_5+\alpha_6)+n_{14235}(-\alpha_6)+n_{14325}(-\alpha_5)\end{aligned}\tag{5.27}$$

by relabeling in the first line and (2.8) at the second line. At another side we have

$$\begin{aligned}\tau_{12345}\to\tau_{12354} &= -\tau_{12345}-\tau_{12435}-\tau_{14235}\\&= n_{12345}(-\alpha_1-\alpha_2-\alpha_4)+n_{12435}(-\alpha_2-\alpha_1-\alpha_3)+n_{13245}(-\alpha_3-\alpha_5-\alpha_6)\\&\quad + n_{13425}(-\alpha_4-\alpha_6-\alpha_5)+n_{14235}(-\alpha_5-\alpha_3-\alpha_1)+n_{14325}(-\alpha_4-\alpha_6-\alpha_2)\end{aligned}\tag{5.28}$$

where in the first line we have used the KK-relation and in the second line we have used the expansion of KK-basis. Comparing above two results, we obtain following equations

$$\begin{aligned}&(-\alpha_1-\alpha_2-\alpha_5)=(-\alpha_1-\alpha_2-\alpha_4),\quad \alpha_2+\alpha_5+\alpha_6=(-\alpha_2-\alpha_1-\alpha_3),\\&(-\alpha_3-\alpha_4-\alpha_6)=(-\alpha_3-\alpha_5-\alpha_6),\quad \alpha_4+\alpha_5+\alpha_6=(-\alpha_4-\alpha_6-\alpha_5)\\&-\alpha_6=(-\alpha_5-\alpha_3-\alpha_1),\quad -\alpha_5=(-\alpha_4-\alpha_6-\alpha_2)\end{aligned}\tag{5.29}$$

Combining (5.26) and (5.29) we can find

$$\alpha_2=\alpha_3=-\frac{2}{5}\alpha_1,\quad \alpha_6=\frac{2}{5}\alpha_1,\quad \alpha_4=\alpha_5=-\frac{1}{5}\alpha_1\tag{5.30}$$

If we put $\alpha_1=\frac{1}{4}$ back, we do reproduce the result (4.14).

Above example can be generalized to any number of elements. The key of our second approach is that if we fully use the potential of relabeling property, we will find expansion of all τ's over BCJ numerator basis $n_{1\sigma n}$ up to an overall factor (for example, α_1 in above example). To determine the overall factor, we need to use relation such as (1.12) in principle. However, based on our examples in section 4, we found following pattern of the expansion of $\tau_{123\ldots(n-1)n}$ over the BCJ numerator basis $n_{1\sigma n}$:

$$\begin{aligned}\tau_{123} &= \frac{1}{2}n_{123}\\ \tau_{1234} &= \frac{1}{3}n_{1234}+\ldots\\ \tau_{12345} &= \frac{1}{4}n_{12345}+\ldots\\ \tau_{123456} &= \frac{1}{5}n_{123456}+\ldots\\ \tau_{1234567} &= \frac{1}{6}n_{1234567}+\ldots\end{aligned}\tag{5.31}$$

We believe this pattern is right although we can not prove it. If we accept this assumption, we do not need to use (1.12) anymore. In other words, with one assumption we can find solutions of τ by using the relabeling property only (cyclic symmetry and KK-relation are still needed). This is the second approach we want to present.

5.3 The implication of permutation $P(1n)$

From our previous discussions, we see that all nontrivial equations for $(n-2)!$ expansion coefficient of $\tau_{123...(n-1)n}$ are given by permutations $P(1n)$ and $P((n-1)n)$. The equations come from permutations $P((n-1)n)$ will be complicated and we do not have a good understanding. However, equations from permutations $P(1n)$ is very simple and we will present them here.

Assuming the expansion

$$\tau_{123..(n-1)n} = \sum_{\sigma\in S_{n-2}} c_\sigma n_{1\sigma n} \tag{5.32}$$

after relabeling $1 \leftrightarrow n$, we have

$$\tau_{n23...(n-1)1} = \sum_{\sigma} c_\sigma n_{n\sigma 1} \tag{5.33}$$

Using the cyclic symmetry and KK-relation we have

$$\tau_{n23...(n-1)1} = \tau_{1n23...(n-1)} = (-)^n \tau_{1(n-1)(n-2)...32n} = (-)^n \sum_{\sigma} c_\sigma n_{1P(\sigma)n} \tag{5.34}$$

where at the last equation we have used the fact that the expansion of $\tau_{1(n-1)(n-2)...32n}$ can be obtained from the expansion of $\tau_{123..(n-1)n}$ by following relabeling

$$P: \quad (2,n-1)(3,n-2)... \tag{5.35}$$

Identifying two different expressions we have

$$(-)^n \sum_{\sigma} c_\sigma n_{1P(\sigma)n} = \sum_{\widetilde{\sigma}} c_{\widetilde{\sigma}} n_{n\widetilde{\sigma}1} = (-)^n \sum_{\widetilde{\sigma}} c_{\widetilde{\sigma}} n_{1(\widetilde{\sigma})^T n} \tag{5.36}$$

thus if

$$P(\sigma) = (\widetilde{\sigma})^T, \qquad \Longrightarrow c_{\widetilde{\sigma}} = c_\sigma \tag{5.37}$$

or more explicitly

$$c_{1\sigma n} = c_{1(P(\sigma))^T n} \tag{5.38}$$

Result (5.38) is the equation coming from permutation $P(1n)$. It is easy to check it with explicit results given in section 4.

From (5.38) we see that there are two special orderings which are singlet under about transformation. They are $c_{1234...(n-1)n}$ and $c_{1(n-1)(n-2)...32n}$. In fact, all coefficients will organize themselves to orbits with one or two elements under the mapping (5.38).

6. Conclusion

In this paper, we have constructed the dual-trace decomposition for Yang-Mills tree amplitudes via kinematic numerators. By imposing cyclic symmetry and KK relation, and the relation between τ's and BCJ numerators in KK-basis, we find solutions of τ's as the linear combinations of BCJ numerators. The dual-trace factors solved in this way satisfy relabeling property. Thus we can get any dual trace factor by relabeling a single τ with given permutation. The relabeling property can be used to fix the dual-trace factors τ.

A. The relabeling of permutation P_{1i}

To complete the proof of relabeling property, we need to prove (5.6) for the permutation $P_{1i} = (1i)$. In this Appendix we will give the detail proof. The equation (5.6) can be rewritten as

$$\begin{aligned}&\sum_{\widetilde{\rho},\widetilde{\sigma}} P(\{i,\rho',1,\sigma',n\}|\{1,\widetilde{\rho},i,\widetilde{\sigma},n\})G(\{1,\widetilde{\rho},i,\widetilde{\sigma},n\}|\{1,\rho,i,\sigma,n\})\\ =&\sum_{\rho'',\sigma''} G(\{i,\rho',1,\sigma',n\}|\{i,\rho'',1,\sigma'',n\})Q(\{i,\rho'',1,\sigma'',n\}|\{1,\rho,i,\sigma,n\}) \qquad (A.1)\end{aligned}$$

The proof can be achieved by the following steps:

Step-1 Finding the explicit expressions of elements in P and Q matrices:

Now let us find explicit expression of elements in P and Q. For permutation $(1i)$ the KK-basis with fixed $(1,n)$ will change to KK-basis with fixed (i,n), thus we need to use various relations, such as Jacobi identity, antisymmetry to get matrix P, and KK relation and cyclic symmetry to get matrix Q.

Let us consider the matrix P first. Using the Jacobi identity and antisymmetry, we have

$$n_{i,\{\rho\},1,\{\sigma\},n} = \sum_{\textit{splittings}\ \rho\to O\{\alpha\}O\{\beta\}} (-1)^{n_\alpha+1} n_{1,\{\alpha\}^T,i,\{\beta\},\{\sigma\},n}. \qquad (A.2)$$

thus elements of matrix P can be read out from the above equation

$$P(i,\{\rho\},1,\{\sigma\},n|1,\{\alpha\}^T,i,\{\beta\},\{\sigma\},n) = \begin{cases} (-1)^{n_\alpha+1} & (\text{if } \rho \text{ can splits into } O\{\alpha\},O\{\beta\}) \\ 0 & \text{Otherwise}\end{cases}. \qquad (A.3)$$

Next we consider the matrix Q. For $\tau_{i,\{\rho\},1,\{\sigma\},n}$ we can use cyclic symmetry and KK relation

$$\tau_{i,\rho,1,\sigma,n} = \tau_{1,\sigma,n,i,\rho} = \sum_{\delta\in OP(\{\sigma\}\bigcup\{\rho^T,i\})} (-1)^{n_\rho+1}\tau_{1,\delta,n}. \qquad (A.4)$$

thus elements of Q can be read out

$$Q(i,\rho,1,\sigma,n|1,\delta,n) = \begin{cases} (-1)^{n_\rho+1} & \text{if } \delta\in OP(\{\sigma\}\bigcup\{\rho^T,i\}) \\ 0 & \text{Otherwise}\end{cases}. \qquad (A.5)$$

Because the ordering $\{\rho^T, i\}$ in (A.5), the nonzero elements of Q must have following form

$$Q(i, \{\alpha^T\}, 1, \{\beta\}, \{\sigma\}, n|1, \{\rho\}, i, \{\sigma\}, n) = (-1)^{n_\alpha+1}, \quad \text{if } \rho \to O\{\alpha\} \bigcup O\{\beta\} \tag{A.6}$$

With explicit expressions of elements of P, the elements of PG in relabeling property (A.1) can be expressed by

$$\sum_{\rho' \to O\{\widetilde{\alpha}\}, O\{\widetilde{\beta}\}} (-1)^{n_{\widetilde{\alpha}}+1} G_{1,n}(1, \{\widetilde{\alpha}^T\}, i, \{\widetilde{\beta}\}, \{\sigma'\}, n|1, \{\rho\}, i, \{\sigma\}, n). \tag{A.7}$$

The elements of GQ in (A.1) is given as

$$\sum_{splittings\ \rho \to O\{\alpha\}, O\{\beta\}} G_{i,n}(i, \{\rho'\}, 1, \{\sigma'\}, n|i, \{\alpha^T\}, 1, \{\beta\}, \{\sigma\}, n)(-1)^{n_\alpha+1}. \tag{A.8}$$

To compare (A.7) and (A.8), using the relabeling invariant property of matrix G, we can exchange the roles of 1 and i in (A.8) and the expression becomes

$$\sum_{splittings\ \rho \to O\{\alpha\}, O\{\beta\}} G_{1,n}(1, \{\rho'\}, i, \{\sigma'\}, n|1, \{\alpha^T\}, i, \{\beta\}, \{\sigma\}, n)(-1)^{n_\alpha+1}. \tag{A.9}$$

Finally the consistency property (A.1) can be rewritten as

$$\begin{aligned} & \sum_{splittings\ \rho' \to O\{\alpha'\}, O\{\beta'\}} (-1)^{n_{\alpha'}+1} G_{1,n}(1, \{\alpha'^T\}, i, \{\beta'\}, \{\sigma'\}, n|1, \{\rho\}, i, \{\sigma\}, n) \\ = & \sum_{splittings\ \rho \to O\{\alpha\}, O\{\beta\}} (-1)^{n_\alpha+1} G_{1,n}(1, \{\rho'\}, i, \{\sigma'\}, n|1, \{\alpha^T\}, i, \{\beta\}, \{\sigma\}, n). \end{aligned} \tag{A.10}$$

It is worth to compare the ordering of parameters of G at both sides. For the first parameter, the two orderings are $\{1, \{\alpha'^T\}, i, \{\beta'\}, \{\sigma'\}, n\}$ vs. $\{1, \{\rho'\}, i, \{\sigma'\}, n\}$ while for the second parameter, the two orderings are $\{1, \{\rho\}, i, \{\sigma\}, n\}$ vs. $\{1, \{\alpha^T\}, i, \{\beta\}, \{\sigma\}, n\}$. Now we need to show sums at the both sides of re-prop2-1 give same number.

Step-2 Further simplification by properties of G

Now let us given a further simplify of the condition (A.1). We notice that $G_{1,n}(\alpha|\beta)$ can be written as

$$G_{1,n}(\sigma'|\sigma) = \sum_{s' \in \{splittings\ \sigma' \to O\{\alpha'\}, O\{\beta'\}\}} \sum_{s \in \{splittings\ \sigma \to O\{\alpha\}, O\{\beta\}\}} \delta(s'|s), \tag{A.11}$$

where $\delta(s'|s)$ is defined as

$$\delta(s'|s) = \begin{cases} 1 & (s' = s) \\ 0 & \text{Otherwise} \end{cases}. \tag{A.12}$$

Substituting (A.11) into (A.10), we get

$$\sum_{\rho'\to O\{\alpha'\},O\{\beta'\}}(-1)^{n_{\alpha'}}\left[\sum_{s',s}\delta(s'|s)\right]=\sum_{\rho\to O\{\alpha\},O\{\beta\}}(-1)^{n_\alpha}\left[\sum_{s',s}\delta(s'|s)\right],\tag{A.13}$$

where we have summed over $s'\in\{\mathcal{S}(\{\alpha'^T\},i,\{\beta'\},\{\sigma'\})\}$ and $s\in\{\mathcal{S}(\{\rho\},i,\{\sigma\})\}$ on the L.H.S, while we have summed over $s'\in\{\mathcal{S}(\{\rho'\},i,\{\sigma'\})\}$ and $s\in\{\mathcal{S}(\{\alpha^T\},i,\{\beta\},\{\sigma\})\}$ on the R.H.S. The above equation can be further rearranged into

$$\begin{aligned}&\sum_{s\in\{\mathcal{S}(\{\rho\},i,\{\sigma\})\}}\left[\sum_{\rho'\to O\{\alpha'\},O\{\beta'\}}(-1)^{n_{\alpha'}}\sum_{s'\in\{\mathcal{S}(\{\alpha'^T\},i,\{\beta'\},\{\sigma'\})\}}\delta(s'|s)\right]\\ &=\sum_{s'\in\{\mathcal{S}(\{\rho'\},i,\{\sigma'\})\}}\left[\sum_{\rho\to O\{\alpha\},O\{\beta\}}(-1)^{n_\alpha}\sum_{s\in\{\mathcal{S}(\{\alpha^T\},i,\{\beta\},\{\sigma\})\}}\delta(s'|s)\right].\end{aligned}\tag{A.14}$$

For a given splitting s the sum in the brackets on the L.H.S. has an useful property

$$\begin{aligned}&\sum_{\rho'\to O\{\alpha'\},O\{\beta'\}}(-1)^{n_{\alpha'}}\sum_{s'\in\{\mathcal{S}(\{\alpha'^T\},i,\{\beta'\},\{\sigma'\})\}}\delta(s'|s)\\ =&\sum_{\rho'\to O\{\alpha'\},O\{\beta'\}}(-1)^{n_{\alpha'}}\sum_{s'\in\{\mathcal{S}(\{\sigma'\}\to\ O\{\sigma'_L\},O\{\sigma'_R\})\}}\delta(O\{\{\alpha'^T\},i,\{\beta'\},\{\sigma'_L\}\},O\{\sigma'_R\}\}|s).\end{aligned}\tag{A.15}$$

For a given splitting s' the sum in the brackets on the R.H.S. has a similar property. The meaning of (A.15) is that for all possible splittings only these with $\{\alpha'^T\},i,\{\beta'\}$ being same subset contribute[3].

Before giving the general proof of the above property (A.15), let us have a look at some examples.

- (1) For the case $n_{\rho'}=1$, i.e., there is only one element in the set ρ', there are two possible splittings: $\{\alpha',\beta'\}=\{\{\ \},\{\rho'_1\}\}/\{\{\rho'_1\},\{\ \}\}$. For the case $\alpha'=\{\rho'_1\}$, the splitting of $\{\rho'_1,i,\sigma'\}$ will have two cases: either ρ'_1,i belong to same subset or ρ'_1,i belongs to different subsets. For the case $\beta'=\{\rho'_1\}$, the splitting of $\{i,\rho'_1,\sigma'\}$ will have two cases: either ρ'_1,i belong to same subset or ρ'_1,i belongs to different subsets. Putting all together, the L.H.S. of (A.15) reads

$$\sum_{\sigma'\to\{\sigma_L\},\{\sigma_R\}}\Bigg[-\delta\left(O\{\rho'_1,i,\sigma'_L\},O\{\sigma'_R\}|s\right)-\delta\left(O\{\rho'_1,\sigma'_L\},O\{i,\sigma'_R\}|s\right)+\delta\left(O\{i,\rho'_1,\sigma'_L\},O\{\sigma'_R\}|s\right)$$

---

[3] One may notice the splitting can be either $O\{\sigma_1,i,\sigma_2\},O\{\sigma_3\}$ or $O\{\sigma_3\},O\{\sigma_1,i,\sigma_2\}$. Since there is no difference for the following discussions between the two kinds of splittings, we just need to deal with only the first kind.

$$+\delta\left(O\{i,\sigma'_L\},O\{\rho'_1,\sigma'_R\}|s\right)\Bigg] \tag{A.16}$$

After summing over all the splittings $\sigma' \to O\{\sigma'\}_L, O\{\sigma'\}_R$, the second term and the fourth term will cancel each other and only two terms are left

$$\sum_{\sigma'\to\{\sigma_L\},\{\sigma_R\}}\Bigg[-\delta\left(O\{\rho'_1,i,\sigma'_L\},O\{\sigma'_R\}|s\right)+\delta\left(O\{i,\rho'_1,\sigma'_L\},O\{\sigma'_R\}|s\right)\Bigg]. \tag{A.17}$$

which is just the R.H.S. of (A.15) when $n_{\rho'}=1$.

- (2) For the case $n_{\rho'}=2$, similar consideration as the case $n_{\rho'}=1$ gives the L.H.S. as

$$\begin{aligned}\sum_{\sigma'\to\{\sigma'_L\},\{\sigma'_R\}}\Bigg\{\Bigg[&\delta\left(O\{\rho'_2,\rho'_1,i,\sigma'_L\},O\{\sigma'_R\}|s\right)+\delta\left(O\{\rho'_2,i,\sigma'_L\},O\{\rho'_1,\sigma'_R\}|s\right)\\
&+\delta\left(O\{\rho'_1,i,\sigma'_L\},O\{\rho'_2,\sigma'_R\}|s\right)+\delta\left(O\{\rho'_2,\rho'_1,\sigma'_L\},O\{i,\sigma'_R\}|s\right)\Bigg]\\
&-\Bigg[\delta\left(O\{\rho'_2,i,\rho'_1,\sigma'_L\},O\{\sigma'_R\}|s\right)+\delta\left(O\{\rho'_2,i,\sigma'_L\},O\{\rho'_1,\sigma'_R\}|s\right)\\
&+\delta\left(O\{\rho'_2,\rho'_1,\sigma'_L\},O\{i,\sigma'_R\}|s\right)+\delta\left(O\{\rho'_1,i,\sigma'_L\},O\{\rho'_2,\sigma'_R\}|s\right)\Bigg]\\
&-\Bigg[\delta\left(O\{\rho'_1,i,\rho'_2,\sigma'_L\},O\{\sigma'_R\}|s\right)+\delta\left(O\{\rho'_1,i,\sigma'_L\},O\{\rho'_2,\sigma'_R\}|s\right)\\
&+\delta\left(O\{\rho'_1,\rho'_2,\sigma'_L\},O\{i,\sigma'_R\}|s\right)+\delta\left(O\{\rho'_2,i,\sigma'_L\},O\{\rho'_1,\sigma'_R\}|s\right)\Bigg]\\
&+\Bigg[\delta\left(O\{i,\rho'_1,\rho'_2,\sigma'_L\},O\{\sigma'_R\}|s\right)+\delta\left(O\{i,\rho'_1,\sigma'_L\},O\{\rho'_2,\sigma'_R\}|s\right)\\
&+\delta\left(O\{i,\rho'_2,\sigma'_L\},O\{\rho'_1,\sigma'_R\}|s\right)+\delta\left(O\{\rho'_1,\rho'_2,\sigma'_L\},O\{i,\sigma'_R\}|s\right)\Bigg]\Bigg\}\end{aligned} \tag{A.18}$$

Again, after summing over all splittings of σ', some terms cancel each other and we are left with

$$\begin{aligned}\sum_{\sigma'\to\{\sigma'_L\},\{\sigma'_R\}}\Bigg[&\delta\left(O\{\rho'_2,\rho'_1,i,\sigma'_L\},O\{\sigma'_R\}|s\right)-\delta\left(O\{\rho'_2,i,\rho'_1,\sigma'_L\},O\{\sigma'_R\}|s\right)\\
&-\delta\left(O\{\rho'_1,i,\rho'_2,\sigma'_L\},O\{\sigma'_R\}|s\right)+\delta\left(O\{i,\rho'_1,\rho'_2,\sigma'_L\},O\{\sigma'_R\}|s\right)\Bigg],\end{aligned} \tag{A.19}$$

which is just the R.H.S. of (A.15) in the case of $n_{\rho'}=2$.

- (3) Similar calculation has be done for the case of $n_{\rho'} = 3$ although we will not present here.

Above examples give the idea of proof. For the case with $n_{\rho'} = r$ after the splitting $\rho' \to O\{\alpha'\}, O\{\beta'\}$ with $n_{\alpha'} = s$ and $n_{\beta'} = r - s$, we need to sum over all splittings of $\{\alpha', i, \beta', \sigma'\}$. In general, the splitting will be $O\{\alpha'_1, i, \beta'_1, \sigma'_1\}, O\{\alpha'_2, \beta'_2, \sigma'_2\}$. Among these two subsets, unlike the subset $O\{\alpha'_1, i, \beta'_1, \sigma'_1\}$ where i distinguish the α' part from β' part, the subset $O\{\alpha'_2, \beta'_2, \sigma'_2\}$ can come from different splitting of $\rho' \to O\{\alpha'\}, O\{\beta'\}$. More explicitly, the last element of α'_2 can be considered as the first element of β''_2. Thus when we put the factor $(-)^{n_{\alpha'}}$ back, the term coming from $n_{\alpha'} = s$ will cancel with the term coming from $n_{\alpha''} = s + 1$. Because this kind of cancelations, only splittings with all elements of $\{\alpha^T\}, i, \{\beta\}$ in a same ordered subset contribute.

Having established (A.15), (A.14) can be rewritten as

$$\begin{aligned}
&\sum_{s\in\{\mathcal{S}(\{\rho\},i,\{\sigma\})\}} \left[\sum_{\rho'\to O\{\alpha'\},O\{\beta'\}} (-1)^{n_{\alpha'}} \sum_{\mathcal{S}(\{\sigma'\}\to\ O\{\sigma'_L\},O\{\sigma'_R\})} \delta(O\{\{\alpha'^T\}, i, \{\beta'\}, \{\sigma'_L\}\}, O\{\sigma'_R\}\}|s) \right] \\
&= \sum_{s'\in\{\mathcal{S}(\{\{\rho'\},i,\{\sigma'\}\})\}} \left[\sum_{\rho\to O\{\alpha\},O\{\beta\}} (-1)^{n_{\alpha}} \sum_{\mathcal{S}(\{\sigma\}\to\ O\{\sigma_L\},O\{\sigma_R\})} \delta(O\{\{\alpha^T\}, i, \{\beta\}, \{\sigma_L\}\}, O\{\sigma_R\}\}|s') \right].
\end{aligned} \tag{A.20}$$

Step-3 Proving the relabeling properties $PG = GQ$ via (A.20)

From step-1 and step-2, we have rewrite the relabeling property $PG = GQ$ to the form (A.20). Now let us prove (A.20) by considering various configurations:

- (1) **Both ρ and ρ' are empty:** In this case, (A.20) becomes

$$\begin{aligned}
&\sum_{\mathcal{S}(\{\sigma\}\to O\{\sigma_L\},O\{\sigma_R\})} \left[\sum_{\mathcal{S}(\{\sigma'\}\to\ O\{\sigma'_L\},O\{\sigma'_R\})} \delta(O\{\sigma'_L\}\}, O\{\sigma'_R\}|O\{\sigma_L\}, O\{\sigma_R\}) \right] \\
&= \sum_{\mathcal{S}(\{\sigma'\}\to O\{\sigma'_L\},O\{\sigma'_R\})} \left[\sum_{\mathcal{S}(\{\sigma\}\to\ O\{\sigma_L\},O\{\sigma_R\})} \delta(O\{\sigma_L\}\}, O\{\sigma_R\}|O\{\sigma'_L\}, O\{\sigma'_R\}) \right],
\end{aligned} \tag{A.21}$$

  which is trivially true.

- (2) **Only one of $\{\rho\}$ and $\{\rho'\}$ is empty:** Assuming $\{\rho\}$ is empty, possible nontrivial terms on L.H.S. of (A.20) is given as

$$\begin{aligned}
&\sum_{\{i,\sigma\}\to O\{i,\sigma_L\},O\{\sigma_R\}} \sum_{\{\sigma'\}\to\ O\{\sigma'_L\},O\{\sigma'_R\}} \delta(O\{i, \rho', \sigma'_L\}, O\{\sigma'_R\}|O\{i, \sigma_L\}, O\{\sigma_R\}) \\
&= \sum_{\{\sigma\}\to O\{\sigma_L\},O\{\sigma_R\}} \sum_{\{\sigma'\}\to\ O\{\sigma'_L\},O\{\sigma'_R\}} \delta(O\{\rho', \sigma'_L\}|O\{\sigma_L\})\delta(O\{\sigma'_R\}|, O\{\sigma_R\}).
\end{aligned} \tag{A.22}$$

where we have used the fact that since the splitting of s with i at the first position, nonzero contributions require the splitting of ρ' to be α' empty.

For the R.H.S. contribution is

$$\begin{aligned}
&\sum_{s'\in\mathcal{S}(\{\rho',i,\sigma'\})}\ \sum_{\{\sigma\}\to\ O\{\sigma_L\},O\{\sigma_R\}} \delta(O\{i,\sigma_L\},O\{\sigma_R\}|s') \\
=&\sum_{\{\sigma'\}\to O\{\sigma'_L\},O\{\sigma'_R\}}\ \sum_{\{\sigma\}\to\ O\{\sigma_L\},O\{\sigma_R\}} \delta(O\{i,\sigma_L\},O\{\sigma_R\}|O\{i,\sigma'_L\},\{\rho',\sigma'_R\}) \\
=&\sum_{\{\sigma\}\to O\{\sigma_L\},O\{\sigma_R\}}\ \sum_{\{\sigma'\}\to\ O\{\sigma'_L\},O\{\sigma'_R\}} \delta(O\{\rho',\sigma'_L\}|O\{\sigma_L\})\delta(O\{\sigma'_R\}|,O\{\sigma_R\}).
\end{aligned} \tag{A.23}$$

where in the second line, to have nonzero result, the splitting of s' is that ρ' and i belong to different subsets. Thus in this case, the relabeling property is also satisfied.

- (3) **Both $\{\rho'\}$ and $\{\rho\}$ are nonempty and they do not share common elements:** Let us consider the L.H.S. of (A.20) first. Since the ρ,ρ' do not share same elements, to have nonzero contribution, we have must α' empty in the splitting of ρ' and ρ,i belong to different subsets in the splitting of s. Thus the L.H.S. of (A.20) is given as

$$\begin{aligned}
&\sum_{\sigma\to O\{\sigma_L\},O\{\sigma_R\}}\ \sum_{\sigma'\to O\{\sigma'_L\},O\{\sigma'_R\}} \delta(O\{i,\rho',\sigma'_L\},O\{\sigma'_R\}|O\{i,\sigma_L\},O\{\rho,\sigma_R\}) \\
=&\sum_{\sigma\to O\{\sigma_L\},O\{\sigma_R\}}\ \sum_{\sigma'\to O\{\sigma'_L\},O\{\sigma'_R\}} \delta(O\{i,\rho',\sigma'_L\}|O\{i,\sigma_L\})\delta(O\{\sigma'_R\}|O\{\rho,\sigma_R\}).
\end{aligned} \tag{A.24}$$

  For the R.H.S., we need the α to be empty in the splitting of ρ and ρ',i belong to different subsets in the splitting of s', thus we get

$$\sum_{\sigma'\to O\{\sigma'_L\},O\{\sigma'_R\}}\ \sum_{\sigma\to O\{\sigma_L\},O\{\sigma_R\}} \delta(O\{i,\rho,\sigma_L\}|O\{i,\sigma'_L\})\delta(O\{\sigma_R\}|O\{\rho,\sigma'_R\}) \tag{A.25}$$

  Thus the relabeling property in this case is satisfied.

- (4) **Both ρ and ρ' are nonempty and they share common elements:** This is a most general case. The L.H.S. of (A.20) is

$$\begin{aligned}
&\sum_{\rho\to O\{\alpha\},O\{\beta\}}\ \sum_{\sigma\to O\{\sigma_L\},O\{\sigma_R\}} \Bigg[\sum_{\rho'\to O\{\alpha'\},O\{\beta'\}} (-1)^{n_{\alpha'}} \\
&\times \sum_{\{\sigma'\}\to\ O\{\sigma'_L\},O\{\sigma'_R\}} \delta(O\{\alpha'^T,i,\beta',\sigma'_L\},O\{\sigma'_R\}\}|O\{\alpha,i,\sigma_L\},O\{\beta,\sigma_R\})\Bigg] \\
=&\sum_{\rho\to O\{\alpha\},O\{\beta\}}\ \sum_{\sigma\to O\{\sigma_L\},O\{\sigma_R\}}\ \sum_{\rho'\to O\{\alpha'\},O\{\beta'\}}\ \sum_{\{\sigma'\}\to\ O\{\sigma'_L\},O\{\sigma'_R\}} \Bigg[(-1)^{n_{\alpha'}}
\end{aligned}$$

$$\times\delta(O\{\alpha'^T\}|\{\alpha\})\delta(O\{\beta',\sigma'_L\}|O\{\sigma_L\})\delta(O\{\sigma'_R\}|O\{\beta,\sigma_R\})\Bigg], \tag{A.26}$$

Similarly, the R.H.S. of (A.20) can be written as

$$\sum_{\rho'\to O\{\alpha'\},O\{\beta'\}}\ \sum_{\sigma'\to O\{\sigma'_L\},O\{\sigma'_R\}}\ \sum_{\rho\to O\{\alpha\},O\{\beta\}}\ \sum_{\{\sigma\}\to\ O\{\sigma_L\},O\{\sigma_R\}}\Bigg[(-1)^{n_\alpha}$$

$$\times\delta(O\{\alpha^T\}|\{\alpha'\})\delta(O\{\beta,\sigma_L\}|O\{\sigma'_L\})\delta(O\{\sigma_R\}|O\{\beta',\sigma'_R\})\Bigg]. \tag{A.27}$$

Changing $\alpha\to\alpha'$ etc, the L.H.S. equals to the R.H.S. in this case.

To sum up, the property (A.20) is satisfied in arbitrary case, then the relabeling condition (A.1) is satisfied for permutation $P(1i)$.

Acknowledgements

Y. J. Du would like to thank Prof. Yong-Shi Wu for helpful suggestions. Y. J. Du is supported in part by the NSF of China Grant No.11105118. CF would like to acknowledge the support from National Science Council, 50 billions project of Ministry of Education and National Center for Theoretical Science, Taiwan, Republic of China as well as the support from S.T. Yau center of National Chiao Tung University. B.F is supported, in part, by fund from Qiu-Shi and Chinese NSF funding under contract No.11031005, No.11135006, No. 11125523.

References

[1] Z. Bern, J. J. M. Carrasco and H. Johansson, "New Relations for Gauge-Theory Amplitudes," Phys. Rev. D **78** (2008) 085011 [arXiv:0805.3993 [hep-ph]].

[2] R. Kleiss and H. Kuijf, "MULTI - GLUON CROSS-SECTIONS AND FIVE JET PRODUCTION AT HADRON COLLIDERS," Nucl. Phys. B **312** (1989) 616.

[3] V. Del Duca, L. J. Dixon and F. Maltoni, "New color decompositions for gauge amplitudes at tree and loop level," Nucl. Phys. B **571** (2000) 51 [hep-ph/9910563].

[4] N. E. J. Bjerrum-Bohr, P. H. Damgaard and P. Vanhove, "Minimal Basis for Gauge Theory Amplitudes," Phys. Rev. Lett. **103** (2009) 161602 [arXiv:0907.1425 [hep-th]].

[5] S. Stieberger, "Open & Closed vs. Pure Open String Disk Amplitudes," arXiv:0907.2211 [hep-th].

[6] S. H. Henry Tye and Y. Zhang, "Dual Identities inside the Gluon and the Graviton Scattering Amplitudes," JHEP **1006** (2010) 071 [Erratum-ibid. **1104** (2011) 114] [arXiv:1003.1732 [hep-th]].

[7] B. Feng, R. Huang and Y. Jia, "Gauge Amplitude Identities by On-shell Recursion Relation in S-matrix Program," Phys. Lett. B **695** (2011) 350 [arXiv:1004.3417 [hep-th]].

[8] Y. Jia, R. Huang and C. -Y. Liu, “$U(1)$-decoupling, KK and BCJ relations in $\mathcal{N} = 4$ SYM,” Phys. Rev. D **82** (2010) 065001 [arXiv:1005.1821 [hep-th]].

[9] H. Tye and Y. Zhang, “Comment on the Identities of the Gluon Tree Amplitudes,” arXiv:1007.0597 [hep-th].

[10] Y. -X. Chen, Y. -J. Du and B. Feng, “A Proof of the Explicit Minimal-basis Expansion of Tree Amplitudes in Gauge Field Theory,” JHEP **1102** (2011) 112 [arXiv:1101.0009 [hep-th]].

[11] H. Kawai, D. Lewellen and H. Tye, ”A Relation Betwwen Tree Amplitudes of Closed and Open Strings”, Nucl.Phys.B269 (1986)1.

[12] N. E. J. Bjerrum-Bohr, P. H. Damgaard, B. Feng and T. Sondergaard, “Gravity and Yang-Mills Amplitude Relations,” Phys. Rev. D **82** (2010) 107702 [arXiv:1005.4367 [hep-th]].

[13] N. E. J. Bjerrum-Bohr, P. H. Damgaard, B. Feng and T. Sondergaard, “New Identities among Gauge Theory Amplitudes,” Phys. Lett. B **691** (2010) 268 [arXiv:1006.3214 [hep-th]].

[14] N. E. J. Bjerrum-Bohr, P. H. Damgaard, B. Feng and T. Sondergaard, “Proof of Gravity and Yang-Mills Amplitude Relations,” JHEP **1009** (2010) 067 [arXiv:1007.3111 [hep-th]].

[15] B. Feng and S. He, “KLT and New Relations for N=8 SUGRA and N=4 SYM,” JHEP **1009** (2010) 043 [arXiv:1007.0055 [hep-th]].

[16] Z. Bern, J. J. M. Carrasco, H. Johansson and , “Perturbative Quantum Gravity as a Double Copy of Gauge Theory,” Phys. Rev. Lett. **105** (2010) 061602 [arXiv:1004.0476 [hep-th]].

[17] Z. Bern, J. J. M. Carrasco, L. J. Dixon, H. Johansson, R. Roiban and , “Simplifying Multiloop Integrands and Ultraviolet Divergences of Gauge Theory and Gravity Amplitudes,” Phys. Rev. D **85** (2012) 105014 [arXiv:1201.5366 [hep-th]].

[18] Z. Bern, C. Boucher-Veronneau, H. Johansson and , “N ¿= 4 Supergravity Amplitudes from Gauge Theory at One Loop,” Phys. Rev. D **84** (2011) 105035 [arXiv:1107.1935 [hep-th]].

[19] C. Boucher-Veronneau, L. J. Dixon and , “N ¿- 4 Supergravity Amplitudes from Gauge Theory at Two Loops,” JHEP **1112** (2011) 046 [arXiv:1110.1132 [hep-th]].

[20] S. G. Naculich, H. Nastase, H. J. Schnitzer and , “Linear relations between N ¿= 4 supergravity and subleading-color SYM amplitudes,” JHEP **1201** (2012) 041 [arXiv:1111.1675 [hep-th]].

[21] S. Oxburgh, C. D. White and , “BCJ duality and the double copy in the soft limit,” JHEP **1302** (2013) 127 [arXiv:1210.1110 [hep-th]].

[22] J. J. M. Carrasco, M. Chiodaroli, M. Gunaydin, R. Roiban and , “One-loop four-point amplitudes in pure and matter-coupled N ¡= 4 supergravity,” JHEP **1303** (2013) 056 [arXiv:1212.1146 [hep-th]].

[23] R. H. Boels, R. S. Isermann, R. Monteiro, D. O’Connell and , “Colour-Kinematics Duality for One-Loop Rational Amplitudes,” arXiv:1301.4165 [hep-th].

[24] N. E. J. Bjerrum-Bohr, T. Dennen, R. Monteiro, D. O’Connell and , “Integrand Oxidation and One-Loop Colour-Dual Numerators in N=4 Gauge Theory,” arXiv:1303.2913 [hep-th].

[25] Z. Bern, S. Davies, T. Dennen, Y. -t. Huang, J. Nohle and , “Color-Kinematics Duality for Pure Yang-Mills and Gravity at One and Two Loops,” arXiv:1303.6605 [hep-th].

[26] C. R. Mafra, O. Schlotterer, S. Stieberger and , “Explicit BCJ Numerators from Pure Spinors,” JHEP **1107** (2011) 092 [arXiv:1104.5224 [hep-th]].

[27] R. Monteiro and D. O’Connell, “The Kinematic Algebra From the Self-Dual Sector,” JHEP **1107** (2011) 007 [arXiv:1105.2565 [hep-th]].

[28] N. E. J. Bjerrum-Bohr, P. H. Damgaard, R. Monteiro and D. O’Connell, “Algebras for Amplitudes,” JHEP **1206** (2012) 061 [arXiv:1203.0944 [hep-th]].

[29] C. -H. Fu, Y. -J. Du and B. Feng, “An algebraic approach to BCJ numerators,” arXiv:1212.6168 [hep-th].

[30] Y. -J. Du, B. Feng and C. -H. Fu, “BCJ Relation of Color Scalar Theory and KLT Relation of Gauge Theory,” JHEP **1108** (2011) 129 [arXiv:1105.3503 [hep-th]].

[31] Z. Bern, T. Dennen, Y. -t. Huang and M. Kiermaier, “Gravity as the Square of Gauge Theory,” Phys. Rev. D **82** (2010) 065003 [arXiv:1004.0693 [hep-th]].

[32] Z. Bern and T. Dennen, “A Color Dual Form for Gauge-Theory Amplitudes,” Phys. Rev. Lett. **107**, 081601 (2011) [arXiv:1103.0312 [hep-th]].